

2010 Breeding Bird Surveys on the Idaho National Laboratory Site

Jericho C. Whiting and Bryan Bybee

April 2011

**Gonzales-Stoller Surveillance
120 Technology Drive
Idaho Falls, ID 83401**

Prepared for:

U.S. Department of Energy-Idaho Operations Office
Environmental Surveillance, Education, and Research Program
Contract No. DE-NE0000-300



EXECUTIVE SUMMARY

Breeding bird surveys (BBSs) have been conducted annually since 1985 (no surveys were conducted in 1992 and 1993) to monitor bird populations on the Idaho National Laboratory (INL) Site. In 2010, we conducted surveys from June 8 to July 1 along 13 established routes, five of which are part of a nationwide survey administered by the U.S. Geological Survey (USGS), and eight of which border INL Site facilities. We documented 5,601 birds from 51 species during these surveys. Bird abundance was greater than the 1985-2009 average of 5,045 birds, but the number of species (i.e., species richness) was lower than the 23-year average of 58.

Compared with past surveys, we observed similar patterns of bird abundance among those species that are typically the most numerous. In 2010, the six species that were surveyed in greatest abundance were the horned lark (*Eremophila alpestris*, $n = 1,447$), western meadowlark (*Sturnella neglecta*, $n = 927$), Brewer's sparrow (*Spizella breweri*, $n = 772$), sage sparrow (*Amphispiza belli*, $n = 520$), Franklin's gull (*Larus pipixcan*, $n = 520$), and sage thrasher (*Oreoscoptes montanus*, $n = 469$). During 24 years of breeding bird surveys on the INL Site, with the exception of the Franklin's gull, these species have been the five most abundant 19 times, and in the remaining five years they were among the six most abundant species. Considering declines reported in populations of sagebrush-obligate species throughout the intermountain west, this trend indicates that the quality of sagebrush-steppe habitat on the INL Site remains stable.

Although three new species were added in the past three years to the list of birds that have been observed at least once during BBS on the INL Site, no new observations were made in 2010. Two species were observed during the surveys that had been recorded in ≤ 3 of the past 23 years. These include yellow warbler (*Dendroica petechia*) and great blue heron (*Ardea herodias*).

Species observed during the 2010 BBS that are considered imperiled or critically imperiled in Idaho include the Franklin's gull ($n = 520$), burrowing owl (*Athene cunicularia*, $n = 3$), greater sage-grouse (*Centrocercus urophasianus*, $n = 1$), and grasshopper sparrow (*Ammodramus savannarum*, $n = 1$).

TABLE OF CONTENTS

EXECUTIVE SUMMARY i

ACRONYMS..... iii

1.0 INTRODUCTION..... 1

 1.1 STUDY AREA 1

 1.2 METHODS..... 3

 1.3 RESULTS AND DISCUSSION..... 5

2.0 SUMMARY 14

 2.1 FUTURE DATA ANALYSIS..... 15

3.0 ACKNOWLEDGEMENTS..... 15

4.0 LITERATURE CITED..... 16

APPENDIX A SUMMARY OF SPECIES BY ROUTE 18

LIST OF FIGURES

Figure 1. Location of Breeding Bird Survey routes on the Idaho National Laboratory Site 2

Figure 2. Number of birds observed during the Breeding Bird Survey on the Idaho National Laboratory Site..... 6

Figure 3. Summary of species assemblage for Breeding Bird Surveys of remote and facility routes on the Idaho National Laboratory Site in 2010 10

Figure 4. Relationship between bird abundance at the Idaho National Laboratory Site and the average June temperature recorded at the Central Facilities Area from 1985 to 2010..... 12

Figure 5. Relationship between bird abundance at the Idaho National Laboratory Site and total June precipitation recorded at the Central Facilities Area from 1985 to 2010..... 13

LIST OF TABLES

Table 1. Number of stops surveyed, species richness, and bird abundance in 2010 for Breeding Bird Survey routes on the Idaho National Laboratory Site 7

Table 2. Summary of species from 13 routes, sorted by abundance, that were observed during the 2010 Breeding Bird Survey on the Idaho National Laboratory Site 8

Table 3. Values for species richness, Shannon Diversity (H) and Equitability (E_H) indices for the 2010 Idaho National Laboratory Site Breeding Bird Surveys..... 14

ACRONYMS

BBS	Breeding Bird Survey
CFA	Central Facilities Area
INL Site	Idaho National Laboratory Site
INTEC	Idaho Nuclear Technology and Engineering Center
MFC	Materials and Fuels Complex
NRF	Naval Reactor Facility
PBF	Power Burst Facility
RTC	Reactor Technology Complex
RWMC	Radioactive Waste Management Complex
TAN	Test Area North
USGS	United States Geological Survey

1.0 INTRODUCTION

The North American Breeding Bird Survey (BBS) was developed by the U.S. Fish and Wildlife Service along with the Canadian Wildlife Service to document trends in bird populations. Pilot surveys began in 1965 and immediately expanded to cover the U.S. and Canada east of the Mississippi and by 1968 included all of North America (Bystrak 1981, Robbins et al. 1986). The BBS program in North America is managed by the U.S. Geological Survey (USGS) and currently consists of over 4,100 routes, with approximately 3,000 of these being sampled each year.

BBS data provide long-term species abundance and distribution trends across a broad-geographic scale. These data have been used to estimate population changes for hundreds of bird species, and they are the primary source for regional conservation programs and modeling efforts (Sauer et al. 2003). Numerous statistical pathways for exploring and analyzing BBS data have been proposed and discussed (James et al. 1996, Link and Sauer 1997, McCulloch et al. 1997, Bart et al. 2004, Sauer et al. 2005). Regardless of differences in opinion concerning the most appropriate analysis techniques, the BBS provides a wealth of information about population trends of birds in North America, and is the foundation for broad conservation assessments extending beyond local jurisdictional boundaries.

The Idaho National Laboratory (INL) Site has five permanent, official BBS routes originally established in 1985 (hereafter referred to as remote routes) and eight additional survey routes near INL Site facilities (hereafter referred to as facility routes). Facility routes were developed to monitor avifauna populations in proximity to anthropogenic activities and disturbances. The annual BBS provides land managers with information regarding the population trends of breeding birds relative to activities conducted on the INL Site. This report summarizes the results from the 2010 BBS and compares species abundance across survey routes with long-term averages.

1.1 STUDY AREA

The INL Site encompasses almost 900 mi² (2,315 km²) on the Upper Snake River Plain in southeast Idaho (Figure 1), and is administered by the U. S. Department of Energy. The INL Site was designated a National Environmental Research Park in 1975 to facilitate research assessing environmental impacts from the development of nuclear energy technologies. This area is located within portions of Bingham, Bonneville, Butte, Clark, and Jefferson Counties. The INL Site has been designated as an Important Bird Area by the Idaho Comprehensive Wildlife Conservation Strategy (Idaho Department of Fish and Game 2005). This designation recognizes wildlife species that are listed by either state or federal agencies and provides a comprehensive listing of the Idaho species of Greatest Conservation Need (Idaho Department of Fish and Game 2005). The INL Site has also been recognized as a Global Important Bird Area by the National Audubon Society.




Figure 1. Location of Breeding Bird Survey routes on the Idaho National Laboratory Site. Blue dots represent survey points along facility routes and red dots represent the same for remote routes.

Topography across the INL Site is mostly flat with an average elevation of 4,985 ft (1,520 m). Other than minor topographic variation created by basalt outcrops, the only significant relief occurs around East and Middle Buttes and the southern portion of the Lemhi Mountains located near the northwest corner of the INL Site.

A description of the climate, geology, and vegetation communities on the INL Site was described in Anderson et al. (1996) and Shive et al. (2011). In general, the INL Site is located in a semi-arid desert that experiences hot, dry summers and cold winters. Annual precipitation on the INL Site averages 8 inches (20 cm), with peak precipitation commonly occurring in spring. The geology is dominated by Quaternary basalt lava flows producing outcrops and lava tubes. Aeolian soils consisting primarily of silt loam and sandy loam are the most common soil type occurring throughout the INL Site, while alluvial soils are more commonly located along the flood plain of the Big Lost River. The INL Site is a shrub-steppe ecosystem dominated by a woody shrub overstory and perennial bunchgrass and forb understory. Wyoming big sagebrush (*Artemisia tridentata wyomingensis*) is the most dominant shrub on the INL Site, while other common species include green rabbitbrush (*Chrysothamnus viscidiflorous*), spiny hopsage (*Grayia spinosa*), shadscale (*Atriplex confertifolia*), winterfat (*Krascheninnikovia lanata*), and

other sagebrush species (*Artemisia* spp.). The most common native grasses are thickspike wheatgrass (*Elymus laceolatus*), bottlebrush squirreltail (*Elymus elymoides*), Indian ricegrass (*Achnatherum hymenoides*), and needle-and-thread grass (*Hesperostipa comata*).

Very little surface water exists during spring and summer on the INL Site. The Big Lost River and Birch Creek drainages are both diverted upstream for agricultural purposes and consequently little, if any, water from these streams reaches the INL Site. During years of high flow, however, water from the Big Lost River can reach the INL Site where it drains into an ephemeral wetland known as the Big Lost River Sinks. This ephemeral wetland provides the only substantial water source for waterfowl and shorebirds on the INL Site, although a number of man-made waste treatment ponds near facilities also provide aquatic habitat for migrating birds.

1.2 METHODS

Data Collection

The BBS is a roadside count of all birds seen or heard along predefined routes. Thirteen BBS routes were surveyed from June 8 to July 1, 2010, consisting of five official USGS BBS routes and eight facility routes that were developed specifically for the INL Site (Figure 1). Each remote survey route is 24.5 miles (39.2 km) with 50 sampling points systematically spaced every 0.5 mile (0.8 km). Facility routes vary in length between 3.6 miles (5.8 km) and 11.9 miles (19.2 km), depending on the size of the facility. Sampling points along facility routes are separated by approximately 0.2 mile (0.4 km).

During surveys, observers followed the North American BBS protocols provided by the USGS Patuxent Wildlife Research Center. At each sampling location (i.e., stop), a trained observer recorded every bird species observed or heard (song) within a quarter-mile radius during a 3-minute interval. Any bird that was suspected of being counted on the previous stop was not recorded again. Additional data such as temperature, wind speed, and sky condition were recorded after every five stops along remote routes, and at the beginning and end of each facility route. Each route was only surveyed when weather conditions were appropriate (e.g., no heavy rain or strong wind). These surveys began one-half hour before sunrise and continued for up to 6 hours until the route was complete. The number of automobiles that passed observers during the 3-minute sampling period was recorded on all remote routes. Also, observers noted whether background noise interfered with audible detection of birds.

Correlation of Bird Abundance and Environmental Factors

In previous reports of BBSs on the INL Site, environmental factors have been investigated to explain variation in observed bird abundance. Between 1985 and 1991, significantly more birds were detected along facility routes in June when the weather was cool and wet than when it was hot and dry (Belthoff et al. 1998). In another report spanning a greater number of years, Belthoff and Ellsworth (1999) showed that high bird abundance in June was significantly correlated with low temperatures and that a non-significant trend existed between high bird abundance and high June precipitation. Interestingly, the removal of an outlier from the 1995 data would have resulted in a statistically significant relationship between abundance and precipitation (Belthoff and Ellsworth 1999). These authors used Spearman rank correlation coefficients to identify whether there was a relationship between bird abundance and June temperature and precipitation (Belthoff and Ellsworth 1999).

The Spearman rank correlation coefficient is a non-parametric test used to investigate the relationship between variables (Spearman 1904). Instead of using the raw abundance data, both variables are ranked in increasing order and the assigned ranks are used in the statistical analyses. The Spearman rank correlation coefficient (r_s) is calculated using the following equation, where (d) is the difference between the ranks and (n) is the sample size.

$$r_s = 1 - \frac{6 \sum d_i^2}{n^3 - n}$$

It is most appropriate to use a different set of equations when there are tied ranks (Thomas 1989), although there is no appreciable difference in the outcome unless there are numerous tied values (Zar 1984). The first equation (see below) is calculated for both variables (x and y) where (t_i) is the number of tied values, and the second equation calculates the Spearman rank correlation coefficient corrected to rank ties (r_s)_c.

$$\sum t_{(xy)} = \frac{\sum (t_i^3 - t_i)}{12}$$

$$(r_s)_c = \frac{(n^3 - n)/6 - \sum d_i^2 - \sum t_x - \sum t_y}{\left[\left[(n^3 - n)/6 - 2 \sum t_x \right] \left[(n^3 - n)/6 - 2 \sum t_y \right] \right]}$$

We used Spearman rank correlation coefficient to investigate relationships between bird abundance and both mean temperature and total precipitation in June since 1985. Weather data were recorded at the Central Facilities Area (CFA) and are available from <http://niwc.noaa.inel.gov/climate.htm>. Statistical significance was calculated using a two-tailed test with $\alpha = 0.05$.

Community Diversity Indices

An ecological community is comprised of all interacting species within a given environment. A community with low species diversity may indicate that an ecosystem is unhealthy or improperly functioning, whereas high species diversity is often used as an indicator of a healthy and stable ecosystem. Consequently, increasing diversity is the goal of many management activities.

Species diversity indices are mathematical methods used to quantify community composition. Many diversity indices are commonly used in ecology and each has particular strengths depending on the data to be analyzed and the questions asked. The simplest estimate of community diversity is species richness, which represents the total number of unique species that occupy an area. Although species richness is a useful measure of diversity, it does not account for differences in abundance between communities. For example, if there are many species for which one individual is observed, richness will be high but may not be comparable with another community that has the same number of species and high abundances of those species. Diversity

indices that consider both species richness and species abundance may provide a more useful measure of community diversity.

Shannon's diversity index (H) is a popular method for quantifying diversity of species in an area (Shannon 1948). This index accounts for both species richness (S) and relative abundance of each species in a community. Shannon's diversity index is derived by first calculating the proportion of species (i) relative to the total number of species (p_i), and then multiplying this proportion by the natural logarithm ($\ln p_i$). The resulting product is then summed across species and multiplied by -1. Shannon's H can range from zero to about 4.6, where higher values represent increasing diversity.

$$H = -\sum_{j=1}^S p_i \ln p_i$$

Another useful measure is Shannon's equitability (E_H). Shannon's equitability represents a measure of evenness, which is how similar species abundance is within a community. E_H ranges from 0 to 1, with 1 representing a completely even community where all species abundances are equal.

$$E_H = H / \ln S$$

Shannon's H and E_H were calculated for all BBS routes, and compared to standard species richness information documented in past reports. We assumed that data obtained from each survey route is an accurate representation of the local bird community.

1.3 RESULTS AND DISCUSSION

Summary Statistics

We documented 5,601 birds during the 2010 survey (Appendix A), which was slightly greater than the average from 1985 to 2009 of 5,045 birds (Figure 2). Species richness of all BBS routes consisted of 51 species, which was lower than the historic average of 58 species (Table 1).

The Tractor Flats route had the highest species richness and Twin Buttes route had the highest bird abundance of all routes (Table 1). Among remote routes, Tractor Flats consistently has had the highest abundance since 1999. The mean bird abundance of this route since 1985 is 710 individuals, which is higher than other remote routes.




Figure 2. Number of birds observed during the Breeding Bird Survey on the Idaho National Laboratory Site. The dashed black line indicates the mean number of birds observed from 1985 to 2010. No BBSs were conducted on the INL Site in 1992 or 1993.

Horned lark (*Eremophila alpestris*) was the most abundant species counted during the 2010 survey with 1,447 individuals representing 26% of all observations (Table 2). This species was observed at 82% (404) of the total stops made during the survey (Table 2). The horned lark is the most abundant species recorded during historic BBSs on the INL Site, and has been the most abundant species annually since 1998.

The six most abundant birds we observed were horned lark ($n = 1,447$), western meadowlark (*Sturnella neglecta*, $n = 927$), Brewer's sparrow (*Spizella breweri*, $n = 772$), sage sparrow (*Amphispiza belli*, $n = 520$), Franklin's gull (*Larus pipixcan*, $n = 520$), and sage thrasher (*Oreoscoptes montanus*, $n = 469$). These six species consisted of > 83% of all observations made in 2010 and they, with the exception of Franklin's gull, were observed on every remote and facility route (Table 2, Appendix A). In the 24 years of INL Site breeding bird surveys, these five species, with the exception of Franklin's gull, have been the most abundant 19 times, and in the remaining five years they were among the six most abundant species. Breeding Bird Surveys in the western U.S. indicate that populations of horned larks, western meadowlarks, Brewer's sparrows, and sage sparrows have all declined across their range (Peterjohn and Sauer 1999). As sagebrush obligates are experiencing population declines from habitat loss and disturbance (Knick et al. 2003), it is encouraging to see a consistently high abundance of these species on the INL Site.

Table 1. Number of stops surveyed, species richness, and bird abundance in 2010 for Breeding Bird Survey routes on the Idaho National Laboratory Site.

Route	Stops	# Species	Abundance
<i>Remote Routes</i>			
Lost River	50	15	468
Circular Butte	50	12	583
Kyle Canyon	50	23	471
Tractor Flats	50	25	805
Twin Buttes	50	22	1138
Subtotal	250	40*	3464
<i>Facility Routes</i>			
CFA	42	22	353
INTEC	25	16	162
MFC	18	18	207
NRF	20	13	237
PBF	28	14	283
RTC	32	15	246
RWMC	20	15	158
TAN	60	13	491
Subtotal	245	37*	2137
Total	495	51	5601

**This value represents the combined number of unique species documented within each route subgroup (i.e., remote vs. facility).*

Rare Observations and Species of Special Concern

Four species were observed during the 2010 BBS that are considered imperiled or critically imperiled in Idaho by the Idaho Department of Fish and Game (2005). These include the burrowing owl (*Athene cunicularia*, $n = 3$), Franklin's gull ($n = 520$), sage-grouse (*Centrocercus urophasianus*, $n = 1$), and grasshopper sparrow (*Ammodramus savannarum*, $n = 1$).

In last year's survey, Franklin's gull numbers were the lowest recorded since 2000, when no birds were observed. This year was the highest since 2002, when there were 764 birds observed. Grasshopper sparrow abundance was at the lowest level since 1996 when no birds were observed. Over the long term, however, these species have been observed more consistently

Table 2. Summary of species from 13 routes, sorted by abundance, that were observed during the 2010 Breeding Bird Survey on the Idaho National Laboratory Site.

Common Name	Scientific Name	n	%	Routes ¹	Stops ²	% ³
Horned Lark	<i>Eremophila alpestris</i>	1,447	25.83	5, 8	404	81.62
Western Meadowlark	<i>Sturnella neglecta</i>	927	16.55	5, 8	311	62.83
Brewer's Sparrow	<i>Spizella breweri</i>	772	13.78	5, 8	360	72.73
Sage Sparrow	<i>Amphispiza belli</i>	520	9.28	5, 8	246	49.70
Franklin's Gull	<i>Larus pipixcan</i>	520	9.28	2, 1	9	1.82
Sage Thrasher	<i>Oreoscoptes montanus</i>	469	8.37	5, 8	295	59.60
Common Raven	<i>Corvus corax</i>	280	5	5, 7	43	8.69
Vesper Sparrow	<i>Pooecetes gramineus</i>	144	2.57	5, 5	84	16.97
Brown-headed Cowbird	<i>Molothrus ater</i>	84	1.5	4, 5	34	6.87
Barn Swallow	<i>Hirundo rustica</i>	60	1.07	1, 7	21	4.24
Mourning Dove	<i>Zenaida macroura</i>	59	1.05	4, 6	39	7.88
Common Nighthawk	<i>Chordeiles minor</i>	51	0.91	5, 7	35	7.07
European Starling	<i>Sturnus vulgaris</i>	40	0.71	1, 5	12	2.42
Black-billed Magpie	<i>Pica pica</i>	23	0.41	2, 0	11	2.22
Brewer's Blackbird	<i>Euphagus cyanocephalus</i>	21	0.37	2, 4	18	3.64
Loggerhead Shrike	<i>Lanius ludovicianus</i>	21	0.37	3, 3	11	2.22
Ferruginous Hawk	<i>Buteo regalis</i>	18	0.32	4, 0	12	2.42
Red-tailed Hawk	<i>Buteo jamaicensis</i>	13	0.23	5, 3	12	2.42
Cliff Swallow	<i>Hirundo pyrrhonota</i>	12	0.21	0, 1	1	0.20
Say's Phoebe	<i>Sayornis saya</i>	12	0.21	0, 5	12	2.42
N. Rough-winged Swallow	<i>Stelgidopteryx serripennis</i>	8	0.14	1, 1	2	0.40
Killdeer	<i>Charadrius vociferus</i>	7	0.12	0, 3	5	1.01
American Kestrel	<i>Falco sparverius</i>	6	0.11	0, 2	5	1.01
Violet-green Swallow	<i>Tachycineta thalassina</i>	6	0.11	1, 1	2	0.40
Gray Flycatcher	<i>Empidonax wrightii</i>	6	0.11	2, 0	5	1.01
Short-eared Owl	<i>Asio flammeus</i>	6	0.11	2, 1	5	1.01
Rock Wren	<i>Salpinctes obsoletus</i>	5	0.09	1, 3	5	1.01
Western Kingbird	<i>Tyrannus verticalis</i>	5	0.09	1, 2	3	0.61
American Robin	<i>Turdus migratorius</i>	4	0.07	1, 1	3	0.61
Mallard	<i>Anas platyrhynchos</i>	4	0.07	0, 2	2	0.40
Northern Harrier	<i>Circus cyaneus</i>	4	0.07	2, 1	4	0.81
Yellow-headed Blackbird	<i>Xanthocephalus xanthocephalus</i>	4	0.07	0, 2	2	0.40
Bank Swallow	<i>Riparia riparia</i>	4	0.07	1, 0	1	0.20
Willet	<i>Catoptrophorus semipalmatus</i>	4	0.07	1, 0	2	0.40
Lazuli Bunting	<i>Passerina amoena</i>	4	0.07	1, 0	2	0.40
Swainson's Hawk	<i>Buteo swainsoni</i>	3	0.05	2, 0	3	0.61
House Sparrow	<i>Passer domesticus</i>	3	0.05	0, 1	1	0.20
Prairie Falcon	<i>Falco mexicanus</i>	3	0.05	2, 1	3	0.61
Blue-gray Gnatcatcher	<i>Poliophtila caerulea</i>	3	0.05	2, 0	3	0.61

Table 2. Continued.

Common Name	Scientific Name	n	%	Routes ¹	Stops ²	% ³
Burrowing Owl	<i>Athene cunicularia</i>	3	0.05	1, 1	2	0.40
Yellow Warbler	<i>Dendroica petechia</i>	3	0.05	1, 1	3	0.61
Red-winged Blackbird	<i>Agelaius phoeniceus</i>	2	0.04	0, 2	2	0.40
Golden Eagle	<i>Aquila chrysaetos</i>	2	0.04	2, 0	2	0.40
Bullock's Oriole	<i>Icterus bullockii</i>	2	0.04	1, 0	2	0.40
House Finch	<i>Carpodacus mexicanus</i>	1	0.02	0, 1	1	0.20
Chipping Sparrow	<i>Spizella passerina</i>	1	0.02	1, 0	1	0.20
Grasshopper Sparrow	<i>Ammodramus savannarum</i>	1	0.02	1, 0	1	0.20
Rock Dove	<i>Columba livia</i>	1	0.02	0, 1	1	0.20
Great Blue Heron	<i>Ardea herodias</i>	1	0.02	0, 1	1	0.20
Greater Sage-Grouse	<i>Centrocercus urophasianus</i>	1	0.02	1, 0	1	0.20
Lark Sparrow	<i>Chondestes grammacus</i>	1	0.02	1, 0	1	0.20

¹The first value represents the number of remote routes at which a species was recorded, and the second value represents the number of facility routes at which a species was recorded.

²Number of stops at which a species was documented.

³Percent of stops (from a total of 495) at which a species was recorded.

during the past 10-15 years than they were during the first decade in which BBS data were collected. Observations of non-imperiled species in 2010 that have rarely been observed during BBSs on the INL Site include yellow warbler (*Dendroica petechia*) and great blue heron (*Ardea herodias*). These two species had been observed in ≤ 3 previous BBSs.

Species Assemblage Summary

Assemblages of bird species in particular habitats within a region provide useful insight about general ecological health of such habitats. For example, if a study area contains large shrubland and grassland habitat patches and the corresponding observations of bird assemblages are low in those areas, this may indicate that the local population is experiencing declines.

The most dominant species assemblage on the INL Site was the shrub-steppe/grassland category, representing nearly 50% of all BBS observations (Figure 3). The shrub-steppe/grassland bird assemblage consistently has the highest bird abundance because the majority of the INL Site consists of shrub-steppe and grassland habitats. The second most abundant species assemblage was sagebrush obligates representing 32% of all observations (Figure 3). Given the regional concern for sagebrush-obligate species (Knick et al. 2003), it is encouraging that they are doing well on the INL Site.

Shrub-steppe/Grassland

Species representing the shrub-steppe/grassland assemblage have always been observed in the greatest number in past BBSs, and they dominated observations in 2010 with 2,785 individuals (49.7%). Common shrub-steppe/grassland species include horned lark, western meadowlark,




Figure 3. Summary of species assemblage for Breeding Bird Surveys of remote and facility routes on the Idaho National Laboratory Site in 2010.

brown-headed cowbird (*Molothrus ater*), and vesper sparrow (*Pooecetes gramineus*). Horned lark ($n = 1,447$) and western meadowlark ($n = 927$) were the most abundant species in this assemblage, and were ranked as the top two most abundant species for the entire survey (Table 2). Annual horned lark observations between 2002 and 2010 have averaged 1,651 birds, whereas the average number observed between 1985 and 2001 was 717 per year. We suspect that the high abundance of horned lark in recent years is a response to wildfires that have converted shrub-dominated habitat into grassland communities. Further investigation of this hypothesis may provide useful insight into the effects of wildfire on bird communities.

Sagebrush Obligates

The sagebrush obligate assemblage had the second highest species abundance with 1,762 individuals (31.5% of total). This assemblage includes Brewer's sparrow, sage sparrow, sage thrasher, and sage-grouse. Brewer's sparrow was the most abundant sagebrush obligate with 772 individuals. These data indicate that populations of sagebrush obligates are thriving on the INL Site. In many other western states, sagebrush obligates are facing significant habitat loss, and consequently sagebrush-obligate species are experiencing population declines (Knick et al. 2003; Sauer et al. 2008). The population trends across the INL Site show a consistently high abundance of sagebrush obligates (Vilord 2007), which is likely because the INL Site is comprised of a large area of relatively undisturbed sagebrush-steppe habitat compared with other areas in the Intermountain West.

Raptors and Corvids

The raptor and corvid assemblage consisted of 382 observations representing 6.8% of the total count. Among these were nine species of raptors (eagles, hawks, falcons, and owls). Ferruginous hawk (*Buteo regalis*) was the most abundant raptor with 18 individuals observed. An observation that was notable in 2010 was that 21 loggerhead shrikes (*Lanius ludovicianus*) were observed. While still lower than the average of 33 loggerhead shrikes per year between

1985 and 2007, it is an increase compared to only 15 loggerhead shrikes observed in 2009 and only 6 observed in 2008.

The corvids include ravens (*Corvus* spp.), crows (*Corvus* spp.), and magpies (*Pica* spp.). The common raven (*C. corax*) was the most abundant species within this assemblage with 280 individuals observed. The number observed in 2010 was the highest abundance since the BBS started. Since egg predation by ravens can negatively impact sage-grouse nest success, it will be important to continue to closely monitor raven abundance, especially if sage-grouse populations continue to decline across the western U.S.

Urbanized and Exotics

The urbanized and exotics assemblage represents birds associated with urban or human-altered environments, which are most commonly found around INL Site facilities. Examples of these species include European starling (*Sturnus vulgaris*), rock dove (*Columba livia*), and American robin (*Turdus migratorius*). This assemblage constituted 2.2% ($n = 121$) of the total observations in 2010. The barn swallow (*Hirundo rustica*) was the most abundant species observed in this assemblage (60 individuals), followed by European starlings (40 individuals).

Waterfowl

Waterfowl are commonly observed during the BBS even though little standing water exists on the INL Site. With the exception of the ephemeral Big Lost River and Sinks wetland, the only standing water bodies on the INL Site are wastewater treatment ponds near facilities. These man-made ponds serve as stopover locations for migrating birds and a number of different species have been observed using these areas since 1985.

We documented four individuals from one waterfowl species, mallards (*Anas platyrhynchos*), representing 0.1% of total observations. As in past years, the mallards were observed from the facility routes.

Shorebirds

We observed 532 individuals representing five species from the shorebird assemblage, which accounted for 9.5% of the total BBS observations. Because standing water is rare on the INL Site, most observations of shorebirds occurred in proximity to waste ponds near facility routes. Franklin's gull ($n = 520$), killdeer (*Charadrius vociferous*, $n = 7$), and willet (*Catoptrophorus semipalmatus*, $n = 4$) comprised nearly all observations. The Franklin's gulls were recorded mainly on the Twin Buttes and Tractor Flats routes. The close proximity of the Tractor Flats route to agricultural areas near Mud Lake is probably why so many gulls were observed.

Other Birds

Four bird species that were not assigned to any species assemblage were observed in 2010. These include the bank swallow (*Riparia riparia*), violet-green swallow (*Tachycineta thalassina*), yellow warbler, and Bullock's oriole (*Icterus bullockii*). Most of these species have been rarely observed during past BBSs; however, from 1985-2003, bank swallows were consistently observed each year (mean = 13). During the past 8 years, bank swallows have been observed during three surveys (2004, 2009, and 2010). Thus, the low number of bank swallow

observations in recent years may indicate that the population is in decline on INL Site survey routes.

Bird Abundance Correlation

Bird abundance was negatively correlated ($r_s = -0.43$, $n = 24$, $P < 0.05$) with mean June temperature (Figure 4). This result supports previous findings from BBSs on the INL Site (Belthoff et al. 1998, Belthoff and Ellsworth 1999), indicating that June temperature should be a consideration when interpreting BBS results. In years where June temperatures are above average, the number of bird observations during the BBS tends to be lower compared with cooler years. The correlation with June temperature and bird abundance thus allows for interpretation of changes in bird abundance across the INL Site, and may help explain annual variability in BBS results.




Figure 4. Relationship between bird abundance at the Idaho National Laboratory Site and the average June temperature recorded at the Central Facilities Area from 1985 to 2010.

Total precipitation in June was not significantly correlated with bird abundance ($r_{sc} = 0.30$, $n = 24$, $P > 0.10$; Figure 5). These results also support previous analyses (Belthoff and Ellsworth 1999). It is interesting that the relationship with June precipitation is not stronger since temperature and precipitation are environmental variables that are inversely related (i.e., in years




Figure 5. Relationship between bird abundance at the Idaho National Laboratory Site and total June precipitation recorded at the Central Facilities Area from 1985 to 2010.

where there is a lot of rainfall, temperatures are typically lower due to evaporative cooling). Although not statistically significant, there is a clear trend towards increased bird abundance as total June precipitation increases. Therefore, precipitation is an important variable to be considered when interpreting changes in annual BBS abundance.

Community Diversity Index

Based on both of Shannon's measures of diversity, the CFA facility route had the most diverse bird community of all 13 routes ($H = 2.53$, $E_H = 0.82$; Table 3), followed by Kyle Canyon ($H = 2.17$, $E_H = 0.69$). Tractor Flats had the highest species richness ($n = 25$). Kyle Canyon had the most diverse bird community among remote routes based on both of Shannon's indicators ($H = 2.17$, $E_H = 0.69$). The TAN route had the lowest diversity among facility routes based on Shannon's measures of diversity ($H = 1.64$; $E_H = 0.64$) and Circular Butte was the least diverse among remote routes based on richness and H . Overall, TAN was the least diverse of all routes, although two routes have slightly lower equitability scores and one has the same equitability score.

Over the past three years, CFA is the only route that has been among the top three in regards to diversity each year. RWMC has been among the three most diverse during three of the past four years. During the same time, Tractor Flats has had the highest or second highest species richness.

Table 3. Values for species richness, Shannon Diversity (H), and Equitability (E_H) indices for the 2010 Idaho National Laboratory Site Breeding Bird Surveys.

Route	Species Richness	Shannon's H	Shannon's E_H
Circular Butte	12	1.71	0.69
Kyle Canyon	23	2.17	0.69
Lost River	15	1.74	0.64
Tractor Flats	25	1.98	0.62
Twin Buttes	22	1.97	0.64
CFA	22	2.53	0.82
INTEC	16	2.09	0.75
MFC	18	1.73	0.60
NRF	13	2.03	0.79
PBF	14	1.76	0.67
RTC	15	1.90	0.70
RWMC	15	2.11	0.78
TAN	13	1.64	0.64

2.0 SUMMARY

As in most previous years, birds belonging to shrub-steppe and grassland community assemblage dominated observations during the 2010 BBS on the INL Site. The total number of birds observed ($n = 5,601$) and species richness ($n = 51$) from all routes was similar to INL Site averages since 1985. Following patterns of abundance from previous BBSs on the INL Site, horned larks were the most abundant species, followed by western meadowlark, Brewer's sparrow, sage sparrow, Franklin's Gull, and sage thrasher. These species, with the exception of the Franklin's Gull, have been consistently among the most abundant species each year of the BBS. These results are important for those concerned about the conservation of sage-steppe ecosystems, because these species are in decline over much of their range. Thus, the habitat quality on the INL Site appears to remain high.

During these surveys we observed 280 ravens. The number observed in 2010 was the highest abundance since the BBS started. Since egg predation by ravens can negatively impact sage-grouse nest success, it will be important to continue to closely monitor raven abundance, especially if sage-grouse populations continue to decline across the western U.S.

Species considered imperiled or critically imperiled in Idaho that were seen during 2010 include: the burrowing owl, Franklin's gull, sage-grouse, and grasshopper sparrow. Other observations of non-imperiled, yet rarely seen birds on the INL Site included yellow warbler and great blue heron.

2.1 FUTURE DATA ANALYSIS

With over two decades of BBS data collected, we are well positioned to conduct a long-term analysis of bird population trends for species occupying the INL Site. Past reports have provided details regarding particular species, but no effort has been made to consider a comprehensive analysis of all BBS data from the INL Site. In the near future, we plan to analyze all data from past BBSs, and to investigate long-term trends in bird abundance and species richness. The results of such an analysis will be submitted to a peer-reviewed journal for publication and will be included in an annual report to the U.S. Department of Energy.

Landscape Change and Habitat Variation

The habitat and vegetation communities across the INL Site are a mosaic of sagebrush-steppe habitat. The INL Site has experienced some large, natural disturbances (e.g., wildfire) which have caused changes in vegetation community composition and distribution across the Site. Little is known, however, concerning responses of bird populations to alterations of habitat composition and distribution across the landscape (Knick and Rotenberry 2002) and how habitat fragmentation can influence local populations. Local bird populations and community assemblages can respond to these habitat changes, and the long-term BBS data should reflect these changes. We will investigate the patterns of habitat change in conjunction with changes in observed bird abundance and richness along routes.

Long-term Community Diversity Trend

Diversity indices have not been calculated each year, and a useful comparison would be to calculate Shannon's H and E_H for all BBS routes for all years to assess which routes have experienced significant change in bird community abundance. The initial community diversity results reported here consider community differences between different routes in the same year. It is unknown how diversity on the same route has changed over time. A number of community similarity indices, such as Morisita's index (Morisita 1959) can be calculated to address this question. We anticipate coupling the results from the spatial analysis described above with the results from community diversity change over time to present a comprehensive description of how bird communities have changed on the INL Site since 1985.

3.0 ACKNOWLEDGEMENTS

We would like to thank Robert Starck for assistance with data collection during spring 2010.

4.0 LITERATURE CITED

- Anderson, J.E., K.T. Ruppel, J.M. Glennon, K.E. Holte, and R.C. Rope. 1996. Plant communities, ethnoecology, and flora of the Idaho National Engineering Laboratory. ESRF-005, Environmental Science and Research Foundation, Idaho Falls.
- Bart, J.B., K.P. Burnham, E.H. Dunn, C.M. Francis, and C.J. Ralph. 2004. Goals and strategies for estimating trends in landbird abundance. *Journal of Wildlife Management* 68: 611-626.
- Belthoff, J.R., L.R. Powers, and T.D. Reynolds. 1998. Breeding birds at the Idaho National Engineering and Environmental Laboratory, 1985 – 1991. *Great Basin Naturalist*. 58:167-183.
- Belthoff, J.R., and E.A. Ellsworth. 1999. Breeding bird surveys at the Idaho National Engineering Laboratory. Unpublished Technical Report, Environmental Science and Research Foundation, Idaho Falls, Idaho.
- Bystrak, D. 1981. The North American Breeding Bird Survey. *Studies in Avian Biology* 6: 34-41.
- Idaho Department of Fish and Game. 2005. Idaho Comprehensive Wildlife Conservation Strategy. Idaho Conservation Data Center, Idaho Department of Fish and Game, Boise, ID. <http://fishandgame.idaho.gov/cms/tech/CDC/cwcs.cfm>
- James, F.C., C.E. McCulloch, and D.A. Wiedenfeld. 1996. New approaches to the analysis of population trends in land birds. *Ecology* 77: 13-27.
- Knick, S.T., D.S. Dobkin, J.T. Rotenberry, M.A. Schroeder, W. Matthew, V. Haegen, and C. Van Riper III. 2003. Teetering on the edge of too late? Conservation and research issues for avifauna of sagebrush habitats. *Condor* 105:611-634.
- Knick, S.T. and J. T. Rotenberry. 2002. Effects of habitat fragmentation on passerine birds breeding in intermountain shrubsteppe. *Studies in Avian Biology* 25: 131-141.
- Link, W.A., and J.R. Sauer. 1997. New approaches to the analysis of population trends in land birds: comment. *Ecology* 78: 2632-2634.
- McCulloch, C.E., F.C. James, and D.A. Wiedenfeld. 1997. New approaches to the analysis of population trends in land birds: reply. *Ecology* 78(8): 2635-2637.
- Morisita, M. 1959. Measuring of the dispersion and analysis of distribution patterns. *Memoirs of the Faculty of Science, Kyushu University, Series E. Biology*. 2: 215-235.
- Peterjohn, B.G. and J.A. Sauer. 1999. Population status of North American grassland birds from the North American Breeding Bird Survey, 1966-1996. *Studies in Avian Biology* 19: 27-44.
- Robbins, C.S., D. Bystrak, and P.H. Geissler. 1986. *The Breeding Bird Survey: its first fifteen years, 1965-1979*. U.S. Fish and Wildlife Service, Resource Publication 157.
- Sauer, J.R., J.E. Fallon, and R. Johnson. 2003. Use of North American Breeding Bird Survey data to estimate population change for bird conservation regions. *Journal of Wildlife Management* 67: 372-389.
- Sauer, J.R., J.E. Hines, and J. Fallon. 2008. *The North American Breeding Bird Survey, results and analysis 1966 – 2007*. Version 5.15.2008. USGS Patuxent Wildlife Research Center, Laurel, MD.
- Sauer, J.R., W.A. Link, J.D. Nichols, and J.A. Royle. 2005. Using the North American Breeding Bird Survey as a tool for conservation: a critique of Bart et al. (2004). *Journal of Wildlife Management* 69(4): 1321-1326.
- Shannon, C.E. 1948. A mathematical theory of communication. *Bell System Technical Journal* 27: 379-423, 623-656.

- Shive, J. P., A. D. Forman, K. Aho, J. R. Hafla, R. D. Blew, and K. T. Edwards. 2011. Vegetation community classification and mapping of the Idaho National Laboratory Site. GSS-ESER-144.
- Spearman, C. 1904. The proof and measurement of association between two things. *American Journal of Psychology* 15: 72-101.
- Thomas, G.E. 1989. A note on correcting for ties with Spearman's ρ . *Journal of Statistical and Computation Simulation* 31: 37-40.
- Vilord, S.J. 2007. Idaho National Laboratory 2006 Breeding Bird Surveys. Unpublished Technical Report, S.M. Stoller Corporation, Idaho Falls, ID.
- Zar, J.H. 1984. *Biostatistical Analysis*. 2nd edition. Prentice Hall. Englewoods Cliffs, New Jersey.

Appendix A

SUMMARY OF SPECIES BY ROUTE 2010

Survey Route: RWMC		
Survey Date: June 11, 2010		
Species	Abundance	Percentage
Brewer's Sparrow	32	20.25
Western Meadowlark	30	18.99
Sage Thrasher	27	17.09
Horned Lark	24	15.19
Barn Swallow	15	9.49
Sage Sparrow	13	8.23
European Starling	5	3.16
Common Nighthawk	3	1.90
Rock Wren	2	1.27
Brewer's Blackbird	2	1.27
Common Raven	1	0.63
Loggerhead Shrike	1	0.63
Red-tailed Hawk	1	0.63
Say's Phoebe	1	0.63
Western Kingbird	1	0.63
Total Individuals		
	158	
Total Species		
	15	

Survey Route: PBF		
Survey Date: June 10, 2010		
Species	Abundance	Percentage
Western Meadowlark	86	30.39
Brewer's Sparrow	78	27.56
Horned Lark	44	15.55
Sage Thrasher	40	14.13
Sage Sparrow	17	6.01
Mourning Dove	4	1.41
Western Kingbird	3	1.06
Short-eared Owl	3	1.06
Common Nighthawk	2	0.71
Say's Phoebe	2	0.71
Common Raven	1	0.35
European Starling	1	0.35
Rock Wren	1	0.35
Loggerhead Shrike	1	0.35
Total Individuals		
	283	
Total Species		
	14	

Survey Route: TAN		
Survey Date: June 29, 2010		
Species	Abundance	Percentage
Horned Lark	194	39.51
Sage Sparrow	116	23.63
Brewer's Sparrow	81	16.50
Sage Thrasher	37	7.54
Vesper Sparrow	37	7.54
Brown-headed Cowbird	11	2.24
Common Raven	5	1.02
Yellow-headed Blackbird	3	0.61
Barn Swallow	2	0.41
Mourning Dove	2	0.41
Western Meadowlark	1	0.20
Rock Dove	1	0.20
Burrowing Owl	1	0.20
Total Individuals		
	491	
Total Species		
	13	

Survey Route: RTC		
Survey Date: June 21, 2010		
Species	Abundance	Percentage
Horned Lark	94	38.21
Brewer's Sparrow	42	17.07
Western Meadowlark	40	16.26
Sage Thrasher	19	7.72
European Starling	14	5.69
Cliff Swallow	12	4.88
Brown-headed Cowbird	6	2.44
Vesper Sparrow	5	2.03
Common Raven	5	2.03
Barn Swallow	4	1.63
Sage Sparrow	1	0.41
Mallard	1	0.41
Prairie Falcon	1	0.41
Red-winged Blackbird	1	0.41
Common Nighthawk	1	0.41
Total Individuals		
	246	
Total Species		
	15	

Survey Route: Circular Butte		
Survey Date: June 22, 2010		
Species	Abundance	Percentage
Horned Lark	166	28.47
Western Meadowlark	145	24.87
Sage Sparrow	95	16.30
Brewer's Sparrow	92	15.78
Sage Thrasher	65	11.15
Brown-headed Cowbird	7	1.20
Common Nighthawk	5	0.86
Common Raven	3	0.51
Vesper Sparrow	2	0.34
Mourning Dove	1	0.17
Red-tailed Hawk	1	0.17
Northern Harrier	1	0.17
Total Individuals		
	583	
Total Species		
	12	

Survey Route: Lost River		
Survey Date: June 24, 2010		
Species	Abundance	Percentage
Horned Lark	193	41.24
Brewer's Sparrow	82	17.52
Western Meadowlark	78	16.67
Sage Sparrow	32	6.84
Sage Thrasher	31	6.62
Vesper Sparrow	28	5.98
Common Raven	11	2.35
Ferruginous Hawk	5	1.07
Red-tailed Hawk	2	0.43
Common Nighthawk	1	0.21
Greater Sage-Grouse	1	0.21
Loggerhead Shrike	1	0.21
American Robin	1	0.21
Brown-headed Cowbird	1	0.21
Swainson's Hawk	1	0.21
Total Individuals		
	468	
Total Species		
	15	

Survey Route: Tractor Flats		
Survey Date: June 14, 2010		
Species	Abundance	Percentage
Horned Lark	241	29.94
Western Meadowlark	199	24.72
Franklin's Gull	133	16.52
Brewer's Sparrow	72	8.94
Sage Thrasher	55	6.83
Sage Sparrow	26	3.23
Mourning Dove	20	2.48
Common Raven	10	1.24
Black-billed Magpie	10	1.24
Vesper Sparrow	9	1.12
Bank Swallow	4	0.50
Willet	4	0.50
Brown-headed Cowbird	3	0.37
Northern Harrier	2	0.25
Short-eared Owl	2	0.25
Brewer's Blackbird	2	0.25
Bullock's Oriole	2	0.25
Burrowing Owl	2	0.25
Ferruginous Hawk	2	0.25
Common Nighthawk	2	0.25
Red-tailed Hawk	1	0.12
Violet-green Swallow	1	0.12
European Starling	1	0.12
Grasshopper Sparrow	1	0.12
Western Kingbird	1	0.12
Total Individuals		
	805	
Total Species		
	25	

Survey Route: Twin Buttes		
Survey Date: June 15, 2010		
Species	Abundance	Percentage
Franklin's Gull	385	33.83
Common Raven	224	19.68
Western Meadowlark	150	13.18
Horned Lark	125	10.98
Brewer's Sparrow	75	6.59
Sage Thrasher	71	6.24
Sage Sparrow	44	3.87
Loggerhead Shrike	12	1.05
Common Nighthawk	10	0.88
Brown-headed Cowbird	8	0.70
Mourning Dove	7	0.62
Vesper Sparrow	7	0.62
Barn Swallow	5	0.44
Red-tailed Hawk	3	0.26
Gray Flycatcher	3	0.26
Ferruginous Hawk	2	0.18
Swainson's Hawk	2	0.18
Prairie Falcon	1	0.09
Blue-gray Gnatcatcher	1	0.09
Brewer's Blackbird	1	0.09
Golden Eagle	1	0.09
Short-eared Owl	1	0.09
Total Individuals		
	1138	
Total Species		
	22	

Survey Route: CFA		
Survey Date: June 28, 2010		
Species	Abundance	Percentage
Horned Lark	68	19.26
Brewer's Sparrow	44	12.46
Brown-headed Cowbird	41	11.61
Western Meadowlark	40	11.33
Sage Sparrow	32	9.07
Sage Thrasher	28	7.93
European Starling	17	4.82
Barn Swallow	16	4.53
Common Raven	14	3.97
Common Nighthawk	14	3.97
Mourning Dove	8	2.27
Brewer's Blackbird	5	1.42
Say's Phoebe	5	1.42
American Kestrel	4	1.13
Vesper Sparrow	4	1.13
American Robin	3	0.85
House Sparrow	3	0.85
Killdeer	2	0.57
Red-tailed Hawk	2	0.57
Great Blue Heron	1	0.28
Loggerhead Shrike	1	0.28
Yellow Warbler	1	0.28
Total Individuals		
	353	
Total Species		
	22	

Survey Route: INTEC		
Survey Date: June 25, 2010		
Species	Abundance	Percentage
Brewer's Sparrow	55	33.95
Horned Lark	27	16.67
Barn Swallow	6	3.70
Sage Sparrow	18	11.11
Sage Thrasher	17	10.49
Western Meadowlark	14	8.64
Common Nighthawk	6	3.70
Violet-green Swallow	5	3.09
Brown-headed Cowbird	4	2.47
American Kestrel	2	1.23
Red-tailed Hawk	2	1.23
Vesper Sparrow	2	1.23
Common Raven	1	0.62
House Finch	1	0.62
Rock Wren	1	0.62
Say's Phoebe	1	0.62
Total Individuals		
	162	
Total Species		
	16	

Survey Route: Kyle Canyon		
Survey Date: June 30, 2010		
Species	Abundance	Percentage
Horned Lark	131	27.87
Sage Sparrow	95	20.21
Sage Thrasher	55	11.70
Brewer's Sparrow	53	11.28
Western Meadowlark	48	10.21
Vesper Sparrow	23	4.89
Mourning Dove	13	2.77
Black-billed Magpie	13	2.77
Ferruginous Hawk	9	1.91
Loggerhead Shrike	5	1.06
Lazuli Bunting	4	0.85
Gray Flycatcher	3	0.64
Common Nighthawk	3	0.64
Common Raven	3	0.64
Northern Rough-winged Swallow	3	0.64
Blue-gray Gnatcatcher	2	0.43
Yellow Warbler	2	0.43
Golden Eagle	1	0.21
Prairie Falcon	1	0.21
Red-tailed Hawk	1	0.21
Lark Sparrow	1	0.21
Chipping Sparrow	1	0.21
Rock Wren	1	0.21
Total Individuals		
	471	
Total Species		
	23	

Survey Route: NRF		
Survey Date: July 1, 2010		
Species	Abundance	Percentage
Horned Lark	67	28.27
Brewer's Sparrow	47	19.83
Sage Sparrow	30	12.66
Vesper Sparrow	27	11.39
Western Meadowlark	20	8.44
Sage Thrasher	20	8.44
Barn Swallow	10	4.22
Northern Rough-winged Swallow	5	2.11
Brown-headed Cowbird	3	1.27
Common Nighthawk	3	1.27
Brewer's Blackbird	2	0.84
Mourning Dove	2	0.84
Killdeer	1	0.42
Total Individuals 237		
Total Species 13		

Survey Route: MFC		
Survey Date: June 8, 2010		
Species	Abundance	Percentage
Western Meadowlark	76	36.71
Horned Lark	73	35.27
Brewer's Sparrow	20	9.66
Brewer's Blackbird	9	4.35
Sage Thrasher	4	1.93
Killdeer	4	1.93
Say's Phoebe	3	1.45
Mallard	3	1.45
European Starling	2	0.97
Franklin's Gull	2	0.97
Mourning Dove	2	0.97
Barn Swallow	2	0.97
Common Raven	2	0.97
Yellow-headed Blackbird	1	0.48
Common Nighthawk	1	0.48
Northern Harrier	1	0.48
Red-winged Blackbird	1	0.48
Sage Sparrow	1	0.48
Total Individuals		
	207	
Total Species		
	18	